



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> A61K 31/00, 47/42, 48/00, 49/00, C07H 21/04, C07K 4/00, G01N 33/566, 33/567	<b>A1</b>	<b>(11) International Publication Number:</b> WO 98/53804 <b>(43) International Publication Date:</b> 3 December 1998 (03.12.98)
<b>(21) International Application Number:</b> PCT/US98/10815 <b>(22) International Filing Date:</b> 28 May 1998 (28.05.98)  <b>(30) Priority Data:</b> 60/047,863 29 May 1997 (29.05.97) US  <b>(71) Applicant (for all designated States except US):</b> AUBURN UNIVERSITY [US/US]; 309 Samford Hall, Auburn, AL 36849-5112 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SMITH, Bruce, F. [US/US]; 624 Woody Drive, Auburn, AL 36832 (US). SAMOILOVA, Tatiana [UA/US]; Apartment 206, 425 Opelika Road, Auburn, AL 36830 (US).  <b>(74) Agents:</b> SPRUILL, W., Murray et al.; Bell Seltzer Intellectual Property Law Group, Alston & Bird LLP, P.O. Drawer 34009, Charlotte, NC 28234 (US).		<b>(81) Designated States:</b> AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> METHODS AND COMPOSITIONS FOR TARGETING COMPOUNDS TO MUSCLE  <b>(57) Abstract</b>  Compositions for use in targeting therapies to muscle cells are provided. The compositions comprise peptides which are capable of binding muscle cells <i>in vivo</i> .		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## METHODS AND COMPOSITIONS FOR TARGETING COMPOUNDS TO MUSCLE

### FIELD OF THE INVENTION

The invention relates to methods for gene therapy, particularly for targeting genes, proteins, pharmaceuticals, or other compounds to muscle.

### BACKGROUND OF THE INVENTION

5           The capabilities to introduce a particular foreign or native gene sequence into a mammal and to control the expression of that gene are of substantial value in the fields of medical and biological research. Such capabilities provide a means for studying gene regulation and for designing a therapeutic basis for the treatment of disease.

10           In addition to introducing the gene into mammals, providing expression of the gene at the site of interest remains a challenge. Methods have been developed to deliver DNA to target cells by capitalizing on indigenous cellular pathways of macromolecular transport. In this regard, gene transfer has been accomplished via the receptor-mediated endocytosis pathway employing  
15   molecular conjugate vectors.

          Inherited diseases of muscle pose a therapeutic challenge, as muscle is the single largest tissue of the body. Pharmacologic approaches do not significantly alter the course of many of these diseases as such approaches fail to correct the underlying genetic deficit. New approaches, relying on the  
20   transfer of genetic material have been advocated. However, current methodologies used for gene therapy are limited in their usefulness with regard to treating myopathies. Local administration of gene therapy vectors or transplantation of donor cells only treat the immediate area of the injection site. Effective therapy with these methods requires multiple injections and carry the  
25   safety risks associated with disease transplantation and the use of immunosuppressive drugs.

          Duchenne muscular dystrophy is probably the most common inherited progressive lethal disorder of mankind. The rate of occurrence of the

-2-

disease is attributable to the very high mutation rate of the gene. The progression of the clinical disease is characterized by skeletal muscle tissue deterioration and wasting. Duchenne muscular dystrophy (DMD) follows a degenerative course which confines sufferers to a wheelchair by the age of about  
5 12 years old and results in death by the third decade due to cardiac or respiratory failure.

DMD results from mutations, mainly frame-shift deletions, in a single, recessively inherited gene. The gene has been cloned and represents the largest gene so far identified in the human genome, spanning at least 2.3  
10 megabases in the short arm of the X chromosome.

Positional cloning of the X-linked gene has revealed that defects of the dystrophin gene lead to either Duchenne or Becker muscular dystrophy (BMD). The 14 kilobase (kb) dystrophin mRNA encodes a 3685 amino acid protein of 427 kilodaltons (kD) with overall similarity to the cytoskeletal  
15 proteins  $\beta$ -spectrin and  $\alpha$ -actinin. These proteins perform structural roles in static and dynamic cellular processes and all cell types. Dystrophin is associated with the cytoplasmic face of the sarcolemma and is thus an essential component of the muscle cytoskeleton. Although the exact function of dystrophin is unknown, it has been postulated to contribute to stabilization of the sarcolemma.

20 The current approaches to the *in vivo* transfer of dystrophin cDNA for the treatment of DMD have involved direct intramuscular injection of naked plasmid DNA or recombinant viral vectors. However, these techniques suffer from the limitation that skeletal muscle comprises a large proportion of the cells of the body, making widespread intramuscular transduction impractical  
25 in a clinical context. Furthermore, some muscles affected in DMD, such as the heart and diaphragm, are not readily accessible to direct injection.

Current approaches to gene therapy in inherited myopathies are limited in practicality either by the mode of administration or the capacity of the vector used for gene transfer. Therefore, methods are needed for targeting  
30 specific compositions to muscle cells.

## SUMMARY OF THE INVENTION

Compositions and methods for targeting genes, proteins, pharmaceuticals, or other compounds to muscle are provided. The compositions comprise peptide sequences which bind muscle, *in vivo*, with high specificity.

5 Both species-specific and species-independent sequences have been determined.

The compositions are useful in gene therapy methods for the treatment of myopathies, gene therapy of disease where muscles potentially serve as reservoirs of protein production, and delivery of a wide variety of drugs to muscles.

## 10 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides the structure of phage in the library.

Figure 2 provides the scheme for selection of tissue-specific phage *in vivo*.

Figure 3 provides the experimental scheme and results.

## 15 DETAILED DESCRIPTION OF THE INVENTION

The invention is drawn to peptide sequences that are capable of binding muscle with high specificity. Both species-specific and species-independent peptide sequences have been identified. The peptide sequences are useful for targeting components to muscle.

20 The peptides of the invention are generally short peptide ligands. The peptide ligands exhibit at least three-fold, preferably ten-fold, more preferably greater than ten-fold binding affinity for muscle cells.

The peptides of the invention are cell-binding and cell-entry peptides. For the most part, the peptides will comprise at least about 5 to about  
25 50 amino acids, preferably at least about 5 to about 30 amino acids, more preferably at least about 7 to about 20 amino acids. It is recognized that consensus sequences may be identified among the peptides that are capable of binding to a target. Such consensus sequences identify key amino acids or patterns of amino acids that are essential for binding. Consensus sequences may  
30 be determined by an analysis of peptide patterns that are capable of binding

muscle cells. Once recognized the consensus regions will be used in constructing other peptides for use in muscular targeting. Such consensus sequences may be tested by constructing peptides and determining the effect of the consensus sequence on binding. In this manner, as long as the consensus  
5 sequence is present, the peptide will bind the target. In some cases, longer peptides will be useful as such peptides may be more easily bound or more readily enter into the target cell.

The peptides can be classified into linear, cyclic and conformational types. While the invention is not bound by any particular mode  
10 of action, it is postulated that the shorter peptides, generally from about 7 to about 20 amino acids are involved in linear binding to the target muscle cells. Longer peptides assume conformational folding and are involved in conformational binding. Cyclic peptide structures can also be constructed for use in the invention. In this manner, a core peptide region such as a consensus  
15 peptide binding sequence will be flanked with identical sequences to form cyclic peptides. Such libraries are available commercially, for example the Ph.D.<sup>TM</sup> Phage display peptide library kits from New England Biolabs, Inc. See also, Parmley *et al.* (1988) *Gene* 73:305-318; Cortese *et al.* (1995) *Curr. Opin. Biotechnol* 6:73-80; Noren (1996) *NEB Transcript* 8(1):1-5; and Devlin *et al.*  
20 (1990) *Science* 249:404-406.

Peptides of the invention can be determined which are capable of binding any type of muscle cells including skeletal, smooth, cardiac, and the like. Based on the selective binding protocols, peptides which are tissue or muscle-type specific or alternatively which are capable of binding to different  
25 muscle cells can be determined. In the same manner, the peptides may be species independent, that is, the peptides will bind to the muscle type from any mammalian species. Alternatively, the peptides may be species specific. By species specific is intended that peptides are specific for muscle cells from a particular species and will not bind to muscle cells from another species.  
30 Therefore, the peptides may be characterized by tissue specificity or alternatively by species specificity. Mammalian species of interest include, but are not limited to human, rat, dog, chimpanzee, etc.

Multiple muscle targets can be utilized to select for muscle-binding peptides. Peptides can be selected against differentiated myotube cell lines from any mammal, as well as against primary muscle samples from mammals, including quadracep, tibialis anterior, soleus, cardiac, and the like.

5           Methods are available in the art for the determination of the peptides of the invention. Such methods include selection from a bacteriophage library which expresses random peptides, mirror image phage display to isolate naturally-occurring L-enantiomers in a peptide library, and the like. See, for example, Barry *et al.* (1996) *Nature Medicine* 2:299-305; Schumacher *et al.* (1996) *Science* 271:1854-1857; Pasqualini *et al.* (1996) *Nature* 380:364-366; and  
10           the references cited therein, herein incorporated by reference.

          Peptides of the invention can be determined from phage libraries which have been used to select random peptides that bind single proteins. See, Barry *et al.* (1996) *Nature Medicine* 2:299-305; Devlin *et al.* (1990) 249:404-  
15           406; Cwirla *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:6378-6382; and the references cited therein. In this manner, peptide libraries can be constructed utilizing a number of random amino acids. The random amino acids are fused to the amino terminus of a phage protein and expressed as a bacteriophage library. See, Barry *et al.* (1996) *Nature Medicine* 2:299-305, herein  
20           incorporated by reference. The phage is hybridized to the cells of interest, at different temperatures, generally about 4°C and about 37°C. After repeated selection, peptides which exhibit a higher affinity for the cells of interest are isolated. Methods for preparing libraries containing diverse populations are also disclosed in Gordon *et al.* (1994) *J. Med. Chem.* 37:1385-1401; Ecker and  
25           Crooke (1995) *BioTechnology* 13:351-360; Goodman and Ro, Peptidomimetics For Drug Design, in "Burger's Medicinal Chemistry and Drug Discovery", Vol. 1, M.E. Wolff (Ed.) John Wiley & Sons 1995, pages 803-861; Blondelle *et al.* (1995) *Trends Anal. Chem.* 14:83-92; and Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, 1989. Each of these  
30           references are herein incorporated by reference.

          After *in vitro* screening of the peptides, the peptides of the invention are selected based on *in vivo* binding. Such methods for *in vivo*

binding are known in the art. See, for example, Pasqualini *et al.* (1996) *Nature* 380:364-366, and the references cited therein. While the invention is discussed in terms of peptides, it is recognized that other molecules may be identified in the same manner. Such molecules include organic chemicals, modified peptides, proteins such as antibodies, antibody fragments, and the like.

The screening method of the invention comprises administering a library of molecules to the muscle cells of interest and identifying those molecules which are capable of binding. See, for example, U.S. Patent No. 5,622,699, herein incorporated by reference.

For peptides capable of binding human muscle cells, chimpanzees can be used as the target mammal. Thus, species independent peptides can be identified which do not cross react with other mammalian species. For peptides which cross-react or are species independent, after screening in a first mammal, the peptides are screened in at least a second mammal and those which bind can be determined. As noted above, because the mammal is typically sacrificed to determine binding, a primate species can be used for human, generally chimpanzees.

The *in vivo* selection method of the invention represents the first methods where peptides have been determined which bind differentiated or "adult" muscle. Previous attempts have selected based on muscle cells in tissue culture. Muscle cells in culture are not characteristic of adult muscle even if collected from developed muscle. Once the cells are cultured they convert to an embryonic phenotype. See Gambke *et al.* (1984) *J. Biol Chem.* 259:12092-12100. Thus, the peptides of the invention have different binding specificities and affinities. It is using the methods of the invention that tissue and species-independent peptides can be determined.

Once peptides have been selected which show an affinity for the target tissue, they may be modified by methods known in the art. Such methods include random mutagenesis, as well as synthesis of the compounds for selected amino acid substitutions. Peptides of various length can be constructed and tested for the effect on binding affinity and specificity. In this manner, the binding affinity may be increased or altered. Thus, peptides may be identified



-7-

which exhibit specific binding to muscle cells, as well as peptides which exhibit specific binding and internalization by the muscle cells of interest.

The peptides find use in targeting genes, proteins, pharmaceuticals, or other compounds to particular muscle tissue. In this  
5 manner, the peptides can be used in any vector systems for delivery of specific nucleotides or compositions to the target cells. By nucleotide is intended gene sequences, DNA, RNA, as well as antisense nucleic acids.

The nucleotide sequences encoding the muscle-specific peptides are also encompassed. Where necessary, the nucleotide sequences can be used  
10 in the construction of fusion proteins or vectors for use in the invention. Such methods are known in the art. Additionally the construction of expression cassettes are known as well as promoters, terminators, enhancers, etc., necessary for expression.

A number of vector systems are known for the introduction of  
15 foreign or native genes into mammalian cells. These include SV40 virus (See, e.g., Okayama *et al.* (1985) *Mol. Cell Biol.* 5:1136-1142); Bovine papilloma virus (See, e.g., DiMaio *et al.* (1982) *Proc. Natl. Acad. Sci. USA* 79:4030-4034); adenovirus (See, e.g., Morin *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:4626; Yifan *et al.* (1995) *Proc. Natl. Acad. Sci. USA* 92:1401-1405; Yang *et al.* (1996)  
20 *Gene Ther.* 3:137-144; Tripathy *et al.* (1996) *Nat. Med.* 2:545-550; Quantin *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:2581-2584; Rosenfeld *et al.* (1991) *Science* 252:431-434; Wagner (1992) *Proc. Natl. Acad. Sci. USA* 89:6099-6103; Curiel *et al.* (1992) *Human Gene Therapy* 3:147-154; Curiel (1991) *Proc. Natl. Acad. Sci. USA* 88:8850-8854; LeGal LaSalle *et al.* (1993) *Science* 259:590-  
25 599; Kass-Eisler *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:11498-11502); adeno-associated virus (See, e.g., Muzyczka *et al.* (1994) *J. Clin. Invest.* 94:1351; Xiao *et al.* (1996) *J. Virol.* 70:8098-8108); herpes simplex virus (See, e.g., Geller *et al.* (1988) *Science* 241:1667; Huard *et al.* (1995) *Gene Therapy* 2:385-392; U.S. Patent No. 5,501,979); retrovirus-based vectors (See, for  
30 example, Curran *et al.* (1982) *J. Virol.* 44:674-682; Gazit *et al.* (1986) *J. Virol.* 60:19-28; Miller, A.D. (1992) *Curr. Top. Microbiol. Immunol.* 158:1-24;

Cavanaugh *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:7071-7075; Smith *et al.* (1990) *Mol. Cell. Biol.* 10:3268-3271; herein incorporated by reference.

In the same manner, the peptides can be used in any mammalian expression vector to target the expression system to the appropriate target muscle  
5 cells. See, for example, Wu *et al.* (1991) *J. Biol. Chem.* 266:14338-14342; Wu and Wu (*J. Biol. Chem.* (1988)) 263:14621-14624; Wu *et al.* (1989) *J. Biol. Chem.* 264:16985-16987; Zenke *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:3655-3659; Wagner *et al.* (1990) 87:3410-3414.

Standard techniques for the construction of the vectors of the  
10 present invention are well-known to those of ordinary skill in the art and can be found in such references as Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd ed. (Cold Spring Harbor, New York, 1989). A variety of strategies are available for ligating fragments of DNA, the choice of which depends on the nature of the termini of the DNA fragments and which choices can be readily  
15 made by those of skill in the art.

Where the peptides of the invention are targeting a gene for expression in muscles, the gene to be expressed will be provided in an expression cassette with the appropriate regulatory elements necessary for expression of the gene. Such regulatory elements are well known in the art and  
20 include promoters, terminators, enhancers, etc.

The peptides of the invention may also be utilized to target liposomes, polylysine, or other polycation conjugates, and synthetic molecules. See, for example, de Haan *et al.* (1996) *Immunology* 89:488-493; Gorlach *et al.* (1996) *DTWDtsch Tierarytl Wochenschr* 103:312-315; Benameur *et al.* (1995)  
25 *J. Phar. Pharmacol.* 47:812-817; Bonanomi *et al.* (1987) *J. Microencapsul* 4:189-200; Zekorn *et al.* (1995) *Transplant Proc.* 27:3362-3363.

In this manner, the peptides of the invention can be used to provide therapies for neuromuscular and muscular diseases, including Duchenne muscular dystrophy, Becker muscular dystrophy, myotonic dystrophy, myotonia  
30 congenita, dermatomyositis, polymyositis, polyneuropathy, and other myopathies. That is, genes, proteins, or pharmaceuticals can be directed to muscles in those patients suffering from the particular disease.

Peptides of the invention also find use in treating cardiac diseases and in targeting drugs to the heart. See, for example, Onodera *et al.* (1997) *Transplant Proc.* 29:1907-1908; Langtry (1997) *Drugs* 53:867-884; Donahue (1997) *Proc. Natl. Acad. Sci. USA* 94:4664-4668; Kass-Eisler *et al.* (1993) *Proc.*  
5 *Natl. Acad. Sci. USA* 90:11498-11502; Guzman *et al.* (1993) *Circ. Res.* 73:1202-1207; Barr *et al.* (1994) *Gene Ther.* 1:51-58; Gojo *et al.* (1997) *J. Ther. Cardiovas. Surg.* 113:10-18; March *et al.* (1995) *Hum Gene. Ther.* 6:41-53; Villarreal *et al.* (1996) *J. Mol. Cell Cardiol.* 28:735-742; Wang *et al.* (1996) *Transplantation* 61:1726-1729; Lee *et al.* (1996) *J. Thorac Cardiovasc. Surg.*  
10 *111:246-252*; and the references cited therein.

In this manner, cardiac peptide directed therapies are useful for the treatment of a number of acquired and inherited cardiovascular diseases. Previous gene transfer approaches have been limited by relatively low efficiencies of gene transduction. Thus, the present approach provides a means  
15 to increase recombinant gene expression and pharmaceutical concentration in the muscular layers of the coronary arteries as well as the myocardium.

The compositions of the invention may be provided as pharmaceutical formulations suitable for parenteral (e.g., subcutaneous, intradermal, intramuscular, intravenous and intraarticular), oral or inhalation  
20 administration. Alternatively, pharmaceutical formulations of the present invention may be suitable for administration to the mucous membranes of the subject (e.g., intranasal administration). The formulations may be conveniently prepared in unit dosage form and may be prepared by any of the methods well-known in the art.

Any inert pharmaceutically-acceptable carrier may be used, such as saline, or phosphate-buffered saline, or any such carrier in which the compositions of the present invention have suitable solubility properties for use in the methods of the present invention. Reference is made to Remington's  
25 *Pharmaceutical Sciences*, Merck Publishing Company Easton, P.A., Osol (ed.) (1980) for methods of formulating pharmaceutical compositions.  
30

The peptides of the invention may also be utilized for gene therapy of disease where muscles may serve as a reservoir of protein production,

-10-

such as diabetes, hemophilia, and the like. Proteins which can be delivered include hGH (Dhawan *et al.* (1991) *Science* 254:1509-1512); factor IX protein (Dai *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:10892-10895; Yas *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:3357-3361; and Roman *et al.* (1992) *Somatic Cell and Molecular Genetics* 18:247-258) insulin; and the like. See, generally, Barr *et al.* (1991) *Science* 245:1507-1509.

In the same manner, the peptides of the invention may be utilized to target pharmaceuticals and chemotherapeutic agents to treat muscle disease such as cancers or tumors of muscle origin. See, for example, Velez-Yanguas *et al.* (1996) *Orthop. Clin. North Am.* 27:545-549; Douglas *et al.* (1995) *Tumor Targeting* 1:67-84; and LeBricon (1995) *Metabolism* 44:1340-1348.

In one example of methods of the invention, the peptides are utilized in a vector system as the ligand-binding domain for the treatment of muscular disease, particularly DMD. The method provides intravenous injection of an expression vector comprising the targeting peptides of the invention operably linked with an expression cassette comprising the Duchenne dystrophy gene to deliver the dystrophin product to the muscles.

Because of the insert size limitations of most retroviral and adenoviral vectors, they may be incapable of accommodating the 14 kb full-length dystrophin cDNA. In these cases, a 6.3 kb dystrophin cDNA may be utilized and expressed in the vector. See, England *et al.* (1990) *Nature* 343:180-182.

Alternatively, adenovirus vectors are known in the art which are capable of expressing the full-length DMD gene. See, for example, Haeker *et al.* (1996) *Human Gene Therapy* 7:1907-1914, which discloses the expression of full-length human dystrophin from an adenoviral vector which has all viral genes deleted. The peptides of the invention could be used to direct such a vector system to skeletal muscle.

The following experiments are offered by way of illustration and not by way of limitation.

## EXPERIMENTAL

Example 1Phage Presentation Libraries May be Utilized to Identify Novel Ligands for Gene Transfer to Muscle

- 5 A key factor in any targeting schema is the availability of appropriate specific molecules on the target cells which can be exploited. Of note in this regard, the available ligands for targeting muscle were until recently limited in number and functional applicability. Many markers of skeletal muscle were described during studies of differentiation and therefore are only expressed
- 10 in myoblasts, mononuclear myocytes, and immature myotubes. However, if skeletal muscle is to be rescued from degeneration due to the absence of dystrophin, mature adult myotubes must be targeted. Thus, the identification of cell surface markers specific to mature myotubes is of critical importance to the implementation of the gene delivery strategies for the treatment of DMD.
- 15 Systems which, while utilizing a variety of different approaches, fundamentally share the similarity of examining libraries of peptides for their ability to bind to specific cell types, both *in vitro* and *in vivo* are useful. See, Figure 1 which sets forth the structure of phage in the library.

Example 2

20 Selection of Peptides against Myoblasts and Myotubes

For MD gene therapy, muscle cells will need to be targeted. Candidate muscle-binding peptides are selected by panning against C<sub>2</sub>C<sub>12</sub> myotubes (Barry *et al.* (1996) *Mature Medicine* 2:299-305).

- In round one of selection, the phage library is injected
- 25 intravenously and multiple tissues may be recovered. In subsequent rounds, the phage purified from specific tissues recovered in round one is injected intravenously, and thus, only that tissue is recovered. In rounds 2-4, separate animals are needed for each tissue being examined. After the last round, the phages are plated on a lawn of bacteria and a number of phage plaques are
- 30 isolated and the DNA sequenced. Of these a certain percentage will represent

-12-

the dominant selected clone. Figure 2 provides a scheme for selection of tissue specific phage.

Therefore, muscle-specific peptides can be identified which will bind to, and be internalized by, the target cells. Specific peptides sequences

- 5 include: WDA NGK T (Trp-Asp-Ala-Asn-Gly-Lys-Thr)  
ASS LNI A (Ala-Ser-Ser-Leu-Asn-Ile-Ala)  
MST QSI N (Met-Ser-Thr-Gln-Ser-Ile-Asn)  
SGL PAY P (Ser-Gly-Leu-Pro-Ala-Tyr-Pro)  
GET RAP L (Gly-Glu-Thr-Arg-Ala-Pro-Leu)  
10 CGH HPV YAC (Cys-Gly-His-His-Pro-Val-Tyr-Ala-Cys)  
QWH GPL H (Gln-Trp-His-Gly-Pro-Leu-His)  
HVH TLP T (His-Val-His-Thr-Leu-Pro-Thr)  
MHH TRF Y (Met-His-His-Thr-Arg-Phe-Tyr)

See, also Table 1 for further *in vivo* selection results.

TABLE 1

SEQUENCE	TISSUE	ANIMAL	NUMBER OF PHAGE CLONES SEQUENCED	FREQUENCY	REMARKS (Selection details)
ASSLNIA	skeletal muscle	mouse	10	40	3 <i>in vitro</i> rounds on C <sub>2</sub> C <sub>12</sub> myotubes (differentiated myoblasts) followed by 2 <i>in vivo</i> rounds on skeletal muscle
WDANGKT	skeletal muscle	mouse	27	19	4 <i>in vivo</i> round sin skeletal muscle
WDANGKT	skeletal muscle	mouse + dog	10	10	2 <i>in vitro</i> on C <sub>2</sub> C <sub>12</sub> myotubes followed by 2 <i>in vivo</i> rounds on skeletal muscle
GETRAPL	skeletal muscle	mouse	27	37	4 <i>in vivo</i> rounds in skeletal muscle
GETRAPL	neuro muscular	mouse	34	12	4 <i>in vivo</i> rounds in neural muscular tissue
CGHHPVYAC	skeletal muscle	dog	20	100	3 <i>in vivo</i> rounds in cardiac muscle. This sequence represents a cyclic peptide where the C residues join to form a ring structure
CGHHPVYAC	cardiac muscle	dog	40	20	3 <i>in vivo</i> rounds in cardiac muscle
SGTLFAN	cardiac muscle	dog	40	28	3 <i>in vivo</i> rounds in cardiac muscle
KAPLTPV	cardiac muscle	dog	40	8	3 <i>in vivo</i> rounds in cardiac muscle
QWHGLPLH	cardiac muscle	mouse	36	20	3 <i>in vivo</i> rounds in cardiac muscle
HVHTLPT	cardiac muscle	mouse	36	11	4 <i>in vivo</i> rounds in cardiac muscle
MHHTRFY	cardiac muscle	mouse	36	8	4 <i>in vivo</i> rounds in cardiac muscle

In the Table, "Four rounds of selection..." means that in round 1, the phage library was injected intravenously, and a specific tissue was recovered. In round 1, multiple tissues are recovered from one animal (skeletal muscle, cardiac muscle, etc.). In any subsequent round, the phage purified from a specific tissue is injected intravenously, and only that tissue is recovered. Thus, in rounds 2-4, separate animals are needed for each tissue being examined. After the last round, the phage are plated on a lawn of bacteria and a number of phage plaques are isolated and the DNA sequenced. This is represented in the Table by "number of phage clones sequenced." Of these, a certain percentage will represent the dominant selected clone. For example, 4 of 10 plaques in the first experiment revealed the ASSLNIA peptide, giving a frequency of 40%.

The experiments disclosed herein, represent the first *in vivo* selection in any species other than mouse. Generally, phage was left to circulate for one hour.

Peptide APVHRPL was selected via two rounds *in vivo*, in canine cardiac muscle. The clone was found in the washed tissue fraction and may represent a very high affinity ligand.

HAIYPRH was selected via four rounds *in vivo*, in murine muscle, at a low frequency (4%). This clone may represent a significant muscle binding peptide.

The frequency of appearance of each amino acid within the library is known. Thus, the frequency of amino acids within the pool of selected peptides can be compared. A number of amino acids have been observed to be over-represented. These amino acids, (H, A, G, V, S, Y, T, N, R) may be significant for muscle binding. Such amino acids may play a role in consensus sequences that are important for binding. One of skill in the art may identify consensus sequences by analysis as well as construction of synthetic sequences and testing binding.



Example 3Selection of A Panel of Muscle-binding Peptides

The peptides identified in Example 2 will be further characterized as described below in Section iii. In addition, a larger panel of muscle-binding peptides will be isolated to increase the likelihood that the most specific and highest affinity muscle-targeting peptides will be identified. Peptides of longer amino acid sequences will also be tested. See Figure 3 for an overall scheme for peptide selection.

i. Selection of Muscle-binding Peptides

Using phage libraries (such as those presented on pIII, 15- and 20-mers presented on pVIII), a variety of myotube-binding peptides will be selected. Multiple muscle targets will be selected against to identify common muscle-binding peptides and increase the likelihood of identifying a wide panel of peptides. Peptides will be selected against differentiated mouse C<sub>2</sub>C<sub>12</sub> and Sol8 myotube cell lines and against primary muscle samples from mouse, rat and dog quadriceps, tibialis anterior, and soleus muscles. A phage selection strategy will be employed to identify peptides which target endocytosing receptors. Selection will be carried through sufficient cycles until increased binding by the population is observed or sequencing of phage clones indicates that the population has collapsed down to a finite number of peptide clones. Clones from this round (round N) will be screened as described below in section iii. Clones from the round N-1 will also be tested, since it is conceivable that truly cell-specific peptides may only transiently appear in the selection process and may be out-competed by less specific, higher affinity/avidity peptides in later rounds of selection. If selection against myotubes does not yield any useful peptides, then selection and screening will be carried out against myoblast cell lines or fresh myoblast isolates.

ii. Selection of Muscle-binding Peptides with New Libraries

While useful peptides have already been isolated from a 20-mer library and a 7-mer library, it should be noted that any one library vastly underrepresents the total number of possible random 20-mer peptides. There is a "space" of  $\sim 10^{26}$  possible 20-mer peptides. Our one library presents  $\sim 10^8$  peptides and therefore represents only 1 in  $10^{18}$  of these possible 20-mers. The number of phage clones is the limitation to how many peptides a library can contain. Therefore, production of more  $10^8$  transformant peptide will produce entirely independent peptides to be screened (up to  $10^{18}$  could in theory be created). It is not known that 20 amino acids are required for cell-binding, however, the 20-mer libraries out complete 12-mer libraries for selection against a variety of cell targets (Barry *et al.* (1996) *Mature Medicine* 2:299-305; and not shown). This suggests large peptides are better for cell binding/entry and that peptides even larger than 20 amino acids may be useful. An additional benefit is derived by using large peptide libraries, since any one 20-mer peptide also represents 14 6-mer peptides, 12 8-mer peptides etc. Given these features, new 20-mer and larger libraries will be created on pIII and pVIII to increase the probability that novel, useful peptides can be identified. New 20-mer libraries will be created using randomly synthesized oligonucleotides (Barry *et al.* (1996) *Mature Medicine* 2:299-305). To create even larger peptide libraries, genomic DNA from *E. coli* or mammalian sources will be randomly sheared to small fragments ( $\sim 200$  base pairs), blunted, and cloned into pIII or pVIII. Once created, these libraries will be selected as in section i and screened as in section iii.

25 iii. Rapid Screening of Peptide Ligands with a Heptapeptide Phage Display Library

a. Ph.D.<sup>TM</sup> Phage Display Peptide Library (New England Biolabs) is a ligand screening system which is based on a combinatorial library of random peptide 7-mers fused to a minor coat protein (pIII) of the filamentous coliphage M13.

-17-

b. The heptapeptide library consists of  $2 \times 10^9$  electroporated sequences amplified once to yield  $\approx 100$  copies of each sequence in 10  $\mu$ l of the supplied phage.

c. Phage display creates a physical linkage between a  
5 selectable function (the displayed peptide sequence) and the DNA encoding that function.

d. This allows rapid identification of peptide ligands for a variety of receptor target molecules by an *in vitro* selection process called biopanning.

10 e. Biopanning is carried out by simply passing the pool of phage displayed peptides over a plate coated with the target, washing away the unbound phage, and eluting the bound phage.

#### Advantages

a. Unlike the use of known ligands, the process requires no  
15 prior knowledge of the biology of the target cells.

b. Molecular recognition and selection are not influenced by the immunogenicity of candidate targets.

c. As a result, peptide ligands should be more easily isolated and incorporated into biological vectors by cloning or by chemical conjugation  
20 to synthetic vectors.

#### iv. Screening Peptides for Specificity

Candidate peptides generated in sections ii and iii will be screened for specific binding to myotubes. Initial screens will compare the binding of these peptides on phage against the myotube lines C<sub>2</sub>C<sub>12</sub> and So18 and against  
25 heterologous cell lines including fibroblasts, macrophages, dendritic cells, and hepatocytes. Peptides will be screened not only for binding muscle cells, but also for low-level binding to hepatocytes (cell lines and primary isolates), since the liver is a tremendous sink for adenovirus *in vivo*. The liver would therefore be a tissue to avoid targeting with new peptides. Likewise, peptides will be  
30 screened for low binding to macrophage and dendritic cells. It is likely that

-18-

delivery of potentially immunogenic gene products and vector antigens to these potent antigen-presenting cells would increase the problem of immune-dependent destruction of adenovirus-transfected cells (Yang *et al.* (1996) *Gene Therapy* 3:412-420).

5                   Some of the current immunogenicity of adenovirus vectors could be due to low-level infection of similar antigen-presenting cells. If so, selected peptides that give lower binding than wild-type fiber protein to these immune cells might actually attenuate the existing immune responses against adenovirus-transfected cells.

10    v.                   Optimization of Peptides by Mutagenesis

As described in section ii, the phage libraries vastly under-represent all possible peptides within a particular library. That any peptides can be selected to bind cells would seem remarkable with only this information. Conversely, this poor representation implies that any peptide selected to bind a  
15   cell has only a  $10^{-18}$  chance of being "the" optimal peptide for binding its target. Therefore, it is likely that any of these lead peptides can be improved by random mutagenesis. Mutagenic libraries will be constructed by insertion of randomly mutated oligonucleotides based on the candidate peptide's sequence into the pIII or pVIII gene in phagemid vector fAFF1 or p8V2 (Barry *et al.*  
20   (1996) *Mature Medicine* 2:299-305). Random insertion of mutant bases in positions 1 and 2 of each codon will be constrained at a ratio of 70:10:10:10 such that 70% of each will contain the normal base and 30% will contain another base. The third position of each codon will be restricted to G or T as described previously (Barry *et al.* (1996) *Mature Medicine* 2:299-305). Once  
25   generated, these libraries will be panned as previously performed (Barry *et al.* (1996) *Mature Medicine* 2:299-305) and consensus amino acids will be identified by sequencing a pool of selected mutant peptide-presenting phage. In addition, the optimal sequence will be determined as described in sections i and iv.

Example 4Vertebrate Animals

Inbred Balb C mice will be used to screen phage presentation libraries. Additional mice will be used following a similar protocol to assay the ability of novel peptide ligands cloned into fiber knob, and into recombinant adenoviruses to bind to muscle. The phage library will be injected intravenously under deep halothane anesthesia. At periods of time ranging from 5 to 30 minutes, the mice will be euthanized by cervical dislocation and immediately immersed in liquid nitrogen. Approximately 20 mice will be allowed to recover from anesthesia following intravenous injection, and euthanized at two days to one week following injection to assess the localization of recombinant adenovirus gene expression.

The initial use of the mouse allows a larger number of potential muscle-specific binding motifs to be screened than would a larger animal model. Initial screens will utilize myoblast and myotube cultures; however, the fetal/neonatal gene expression pattern of these cells while in culture may not permit the isolation of binding moieties appropriate for adult skeletal muscle. Thus, the use of *in vivo* screening of phage libraries will significantly shorten the interval involved in discovery of these novel binding moieties. Subsequently, the ability of modified fiber, and then of modified adenovirus to specifically target muscle after intravenous injection can only be tested *in vivo*. The use of the M-PFK deficient dog model provides a readily assessable animal model of a myopathy in which the ability of tropism modified adenovirus to deliver a therapeutic gene can be investigated. (See Example 5).

Example 5Therapy in a Canine Modeli. Pre-treatment Studies

Neonatal dogs (1 day to 1 week old) from an M-PFK colony will be tested for M-PFK deficiency using the PCR-based allele-specific test that we devised. Five juvenile (4-9 months) dogs, which have been determined to be

-20-

homozygous mutant for M-PFK deficiency, will be used in this study. Each dog will have skeletal muscle biopsies taken from the gastrocnemius and triceps muscles. Portions of these biopsies will be frozen and evaluated histochemically. Tissue samples will also be homogenized and assayed for M-PFK activity.

- 5 Detailed medical records are kept on all colony animals, and any episodes of anemia or pigmenturia will be recorded. Pre-treatment magnetic resonance spectroscopy (MRS) studies will be performed on each animal to establish parameters for these investigations. Whole blood and muscle will be frozen at -70°C to provide pre-treatment controls for Southern hybridizations, PCR
- 10 amplification of vector sequences, PFK activity studies, and M-PFK-specific immunoblots. Serum samples will be frozen as controls for the antigenicity studies described below.

ii. In vivo Gene Delivery to M-PFK Deficient Dogs

- After completion of the pre-treatment studies, each dog will be
- 15 sedated, and a 21 gauge intravenous catheter placed in the cephalic vein. Tropism-modified vectors comprising the peptides and expression cassettes containing M-PFK will be delivered as a slow injection. During injection, the dogs will be monitored for signs associated with adverse reactions such as anaphylaxis, and should such reactions occur, administration of the vector will
- 20 be suspended and appropriate treatments initiated.

iii. Studies to Detect the Transferred Canine M-PFK cDNA

- Skeletal muscle biopsies will be taken from the contralateral gastrocnemius and triceps muscle at one, two, three, four, six and eight weeks following the injection of the vector, and monthly thereafter. DNA will be
- 25 recovered from this tissue sample and examined by PCR for the transferred virus.

iv. Immunological Detection of Tissue Specific Expression of M-PFK

To determine if the vector which has been transferred to the muscle is expressing M-PFK, M-PFK specific immunoblots will be performed. One hundred to one hundred fifty micrograms of protein are loaded into each well of a 7.5% polyacrylamide gel. Following electrophoresis the gel is stained with coomassie brilliant blue to show protein content, and the electrophoretically transferred to 0.45 micron nitrocellulose membranes. The membranes are blocked with 5% milk powder for 1 hr, followed by 0.05% Tween 20 for 30 min. The membranes are then incubated with the primary antibody, guinea pig anti-rabbit M-PFK, for 18 hr at room temperature, with gentle agitation. Three washes with Tris-buffered saline (0.2 M Tris, 5 M NaCl, pH 7.5; TBS), 30 seconds to one minute each, are followed by a 3 hr incubation with peroxidase-conjugated rabbit anti-guinea pig IgG (Sigma), at room temperature. A final three rinses in TBS are followed by the addition of 4 chloro-naphthol to allow for color development.

v. Biochemical Detection of Enzyme Activity from Transferred M-PFK

To determine if the expressed M-PFK possesses enzymatic activity, the muscle biopsy samples will be evaluated by biochemical assay. Phosphofructokinase from muscle biopsies will be prepared for activity assay by homogenization for 15 seconds in 5 volumes of PFK extraction buffer [50 mM Tris-phosphate, pH 8.0, 25 mM NaF, 0.1 mM ethylenediamine-tetraacetic acid (EDTA), 25 mM  $(\text{NH}_4)_2\text{SO}_4$ , 1 mM adenosine triphosphate (ATP), 10 mM dithiothreitol (DTT), 0.5 mM phenylmethyl-sulfonyl fluoride (PMSF), 1 mM p-aminobenzamidine, 250  $\mu\text{g/liter}$  Leupeptin, 0.1 mM Na-p-tosyl-L-lysine chloromethyl ketone]. The homogenate is then sonicated and centrifuged at 20,000 x g for 1 hour. The supernatant is saved and stored at -20°C. Enzyme activity will be determined by a calorimetric reaction in 50 mM tris(hydroxymethyl)methylglycine, pH 8.4, 5mM  $\text{MgCl}_2$ , 0.15 mM NADH, 0.1 mM DTT, 0.25 mM ATP, 2.4 mM Fructose-6-phosphate, 0.18 units aldolase, 0.6 units triose phosphate isomerase, and 0.1 unit glycerophosphate

-22-

dehydrogenase. The decrease in absorbance at a wavelength of 340 nm is monitored for 10 min at 26°C and one unit of PFK activity is defined as the amount of enzyme needed to convert 1  $\mu$ mole of fructose-6-phosphate to fructose-1, 6-diphosphate in one minute. All studies will be compared to pretreatment values as well as values from ag-matched normal and M-PFK deficient dogs, and dogs treated with a control vector.

vi. Histochemical Localization of Tissue-specific Expression of M-PFK

- Determination of the number and type of muscle cells expressing M-PFK activity will be performed by histochemical staining for M-PFK. Briefly, small pieces of muscle, frozen in liquid nitrogen-cooled isopentane, will be mounted in a cryostat and sectioned at 6 microns. These sections will be placed on microscope slides, dried, and incubated in a solution containing 10 mM  $\text{Na}_2\text{HAsO}_4$ , 2 mM fructose-6 phosphate, 1 mM diphosphopyridine nucleotide (DPNH) 1 mM ATP, 1 mM  $\text{MgSO}_4$ , and 4 mg/ml NBT. The present of PFK activity is seen as a dark blue stain within the cells.

vii. Evaluation of the Efficacy of Gene Transfer in Treating M-PFK Deficiency

- To evaluate the suitability of the molecular conjugate vectors for treating inherited myopathies, the effects of treatment on the clinical signs of M-PFK deficiency will be evaluated. The dogs will be observed closely for signs of myopathy, including fatigue, exercise intolerance, and muscle cramping. Light levels of long duration exercise such as walking will be used to attempt to induce signs of myopathy. If stable expression is seen in muscle biopsy specimens exceeding one month post injection, MRS studies will be performed to determine the effect of M-PFK expression on the metabolic block. With the dog placed under general anesthesia, the muscle will be tetanically stimulated and the level of inorganic phosphorus determined. The results of these studies will be compared to the pretreatment values to determine the extent of



improvement in muscle function. Additionally, the dogs will be closely observed during this procedure for episodes of cramping or muscle contracture.

viii. Detection of Immune Response to the Adenoviral Vector

To determine if the vectors induce an immune response in the animal being treated, serum samples taken prior to treatment and each biopsy will be hybridized with nitrocellulose blots of adenoviral proteins run on SDS polyacrylamide gels. A secondary, labeled anti canine IgG will be utilized to visualize any canine anti-adenoviral antibodies. Antibodies to the transferred M-PFK gene are not expected as the canine M-PFK will be used and the deficiency is due to a point mutation.

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

## THAT WHICH IS CLAIMED:

1. A method for determining muscle-specific peptides, said method comprising constructing a bacteriophage library which expresses random peptides at the amino terminus of a phage protein, and hybridizing the resulting  
5 phage to muscle cells of interest wherein said peptides are selected based on in vivo binding.
2. The method of claim 1, wherein said peptides are random peptides.
3. The method of claim 1, wherein said peptides are 7-mer  
10 peptides.
4. The method of claim 1, wherein said muscle-specific peptides are species specific.
5. The peptides of claim 1, wherein said muscle-specific peptides are species independent.
- 15 6. The method of claim 1, wherein said muscle-specific peptides are capable of binding to specific muscle cells selected from the group consisting of smooth, skeletal and cardiac muscle cells.
7. The method of claim 6, wherein said muscle cells are skeletal muscle cells.
- 20 8. The method of claim 6, wherein said muscle cells are cardiac muscle cells.

-25-

9. A method for targeting a pharmaceutical composition to muscle cells, said method comprising ligating a muscle-specific peptide to said pharmaceutical composition.

10. The method of claim 9, wherein said pharmaceutical  
5 composition is selected from drugs, proteins, and nucleic acids.

11. The method of claim 9, wherein said pharmaceutical composition is useful for treating a myopathy.

12. The method of claim 9, wherein said peptides are random peptides.

10 13. The method of claim 9, wherein said peptides are 7-mer peptides.

14. The method of claim 9, wherein said muscle-specific peptides are species specific.

15 15 peptides are species independent.

16. The method of claim 9, wherein said muscle-specific peptides are capable of binding to specific muscle cells selected from the group consisting of smooth, skeletal and cardiac.

17. The method of claim 16, wherein said muscle cells are  
20 skeletal muscle cells.

18. The method of claim 16, wherein said muscle cells are cardiac muscle cells.

-26-

19. An isolated muscle-specific peptide which is capable of specific binding to muscle cells *in vivo*.

20. The peptide of claim 19, wherein said peptide comprises about 7 to about 20 amino acids.

5           21. The peptide of claim 19, wherein said muscle cells are selected from smooth, cardiac, or skeletal muscle cells.

22. The peptide of claim 21, wherein said muscle cells are skeletal muscle cells.

10           23. The peptide of claim 21, wherein said muscle cells are cardiac.

24. The peptide of claim 19, wherein said peptide is species specific.

25. The peptide of claim 19, wherein said peptide is species independent.

15           26. A nucleotide sequence encoding the peptide of any of claims 19-25.

27. A method for treating a patient having a muscular disorder, said method comprising administering to said patient a pharmaceutically-effective amount of a composition comprising a muscle-specific peptide operably linked to a pharmaceutical composition useful for  
20           treating said disorder.

28. The method of claim 27, wherein said peptide is species specific.

-27-

29. The method of claim 28, wherein said peptide is species independent.

30. The method of claim 27, wherein said peptide specifically binds cardiac muscle.

5 31. The method of claim 27, wherein said peptide specifically binds skeletal muscle.

32. The method of claim 27, wherein said peptide specifically binds smooth muscle.